

Research Paper

# Study on Genetic Parameters for Kleiber Ratio in Sonadi Sheep

Gaurav Kumar Bansal<sup>1\*</sup>, Krishna Nand Bansal<sup>2</sup>, Vishnu Kumar<sup>1</sup>, R.K. Nagda<sup>1</sup>,  
Renuka Hada<sup>1</sup> and Ravta Ram<sup>3</sup>

<sup>1</sup>Department of Animal Genetics and Breeding, College of Veterinary and Animal Science, Navania, Udaipur, Rajasthan, India

<sup>2</sup>Department of Veterinary Gynaecology and Obstetrics, College of Veterinary and Animal Science, Navania, Udaipur, Rajasthan, India

<sup>3</sup>SRE, Krishi Vigyan Kendra, Gudamalani, Barmer, Rajasthan, India

\*Corresponding author: grvbns1321@gmail.com

Received: 13-02-2023

Revised: 29-05-2023

Accepted: 07-06-2023

## ABSTRACT

Data of 1408 Sonadi lambs born during 2012-2019 under Mega Sheep Seed Project at CVAS Navania Udaipur (Rajasthan) were analyzed to assess the influence of period, season, sex and type of birth on Kleiber ratio. The overall least-squares means along with standard error of Kleiber ratio - KR1 (0-3M), KR2 (3-6M), KR3 (6-9M) and KR4 (9-12M) were 13.16±0.30, 6.59±0.33, 4.04±0.31 and 5.24±0.35, respectively. Effect of sire was highly significant ( $P \leq 0.01$ ) on KR1, KR2 and KR3 whereas significant ( $P \leq 0.05$ ) on KR4. Period of birth had highly significant ( $P \leq 0.01$ ) effect on all the KRs. Season of birth had highly significant ( $P \leq 0.01$ ) effect on KR4 only whereas sex of lamb had highly significant ( $P \leq 0.01$ ) effect on KR2 only. Type of birth had significant ( $P \leq 0.05$ ) effect on KR4 only. Estimates of heritability for KR1, KR2, KR3 and KR4 were 0.82±0.18, 0.38±0.11, 0.51±0.15 and 0.23±0.13, respectively. Genetic correlation was negative between KR1 and other KRs whereas it was positive among other KRs. The estimate of phenotypic correlation among Kleiber ratios was negative and low to high.

**Keywords:** Kleiber ratio, Sonadi sheep, least-square mean, heritability, genetic and phenotypic correlation

Sheep (*Ovis aries*) is one of the important species of livestock contributing to meat and wool production. Sheep are capable of living in all kind of environment and thrive well in hot and arid regions. India ranks 3<sup>rd</sup> in sheep population having a number of 74.26 Million and contributes for 13.8% of India's total livestock population (Livestock census, 2019). Rajasthan with 7.9 million of sheep population is the 4<sup>th</sup> largest sheep (10.63% of total sheep in India) rearing state of the India.

Sonadi sheep is an important mutton breed of southern Rajasthan and prevails in southern sub-humid zone comprising Udaipur, Chittorgarh, Rajsamand, Dungarpur, Banswara districts of Rajasthan. It is generally reared under semi extensive system. It has the unique characteristic of golden fibre and survival on scarce fodder condition during drought. The fleece of Sonadi

sheep is extremely coarse and hairy. Belly and legs are devoid of wool. Sonadi sheep has good potential for meat (13.72 kg at slaughter) and wool (0.5-1.2 kg/annum) production.

Kleiber ratio (KR), defined as growth rate divided by metabolic body weight ( $W^{0.75}$ ), was suggested for growth efficiency (Kleiber, 1947). The relation of growth rate to metabolic weight or Kleiber ratio (KR) was developed as an alternative ratio to address problem in rangeland animals (Arthur *et al.* 2001). In order to increase the efficiency of feed conversion, use of Kleiber ratio (KR) in selection

**How to cite this article:** Bansal, G.K., Bansal, K.N., Kumar, V., Nagda, R.K., Hada, R. and Ram, R. (2023). Study on Genetic Parameters for Kleiber Ratio in Sonadi Sheep. *Theriogenology Insight: An Int. J. Reprod. of Anim.*, 13(01): 19-23.

Source of Support: None; Conflict of Interest: None



programs is very essential (Eskandarinasab *et al.* 2010). The greater correlation between Kleiber ratio and feed efficiency indicates that animals with higher Kleiber ratio require lower energy for their maintenance (Kumar *et al.* 2017).

## MATERIALS AND METHODS

Data for the investigation were collected from the farm unit of Sonadi sheep maintained under Mega Sheep Seed Project (MSSP) at Vallabh Nagar, Udaipur (Rajasthan). A total record of 1408 lambs of Sonadi sheep having 718 females and 690 males spread over period of 8 years from 2012 to 2019 comprised the material for this study. The breeding tract of Sonadi sheep falls under southern sub-humid plain zone of Rajasthan. The Sonadi Sheep are being maintained under semi-intensive management system. Sheep are remained on pasture everyday for 6-8 hours for grazing. The lambs are weaned at 3 months of age. Kleiber ratio is ratio of Average Daily Gain (ADG) in gm and Metabolic Body Weight ( $W^{0.75}$ ) in Kg. Kleiber ratio KR1 (0-3 months), KR2 (3-6 months), KR3 (6-9 months) and KR4 (9-12 months) were calculated by following formula:-

$$\text{Kleiber Ratio (KR)} = \text{ADG} / W^{0.75}$$

Where,

ADG = Average daily gain for the period expressed in g/day

$W^{0.75}$  = Metabolic body weight in Kg. at the older age of the period for which KR is calculated.

Year of birth were grouped into four periods as P1 (2012-2013), P2 (2014-2015), P3 (2016-2017) and P4 (2018-2019). Season of birth was classified into three seasons as S1 (monsoon), S2 (winter) and S3 (summer). Sex of lamb was classified into male and female whereas type of birth into single and twins. The data on Kleiber ratio was analyzed through Mixed Model Least-Squares and Maximum Likelihood method designed by Harvey (Harvey, 1990). Effect of various non-genetic factors on Kleiber ratio was estimated through the following statistical model:

$$Y_{ijklmn} = \mu + A_i + B_j + C_k + D_l + E_m + e_{ijklmn}$$

Where,

$Y_{ijklmn}$  = Growth record of the  $n^{\text{th}}$  progeny of  $i^{\text{th}}$  sire belonging to  $l^{\text{th}}$  sex,  $k^{\text{th}}$  season,  $j^{\text{th}}$  period and  $m^{\text{th}}$  type of birth

$\mu$  = Population mean

$A_i$  = Random effect of  $i^{\text{th}}$  sire

$B_j$  = Fixed effect of  $j^{\text{th}}$  period of birth ( $i = 1, 2, 3, 4$ )

$C_k$  = Fixed effect of  $k^{\text{th}}$  season of birth ( $j = 1, 2, 3$ )

$D_l$  = Fixed effect of  $l^{\text{th}}$  sex of lamb ( $k = 1, 2$ )

$E_m$  = Fixed effect of  $m^{\text{th}}$  type of birth ( $l = 1, 2$ )

$e_{ijklmn}$  = Residual error, NID ( $0, \sigma^2$ )

Duncan's Multiple Range Test (DMRT) as modified by Kramer was used to make pair wise comparison among the least squares means (Kramer, 1956).

## RESULTS AND DISCUSSION

The least-squares means for KR1, KR2, KR3 and KR4 were  $13.16 \pm 0.30$ ,  $6.59 \pm 0.33$ ,  $4.04 \pm 0.31$  and  $5.24 \pm 0.35$ , respectively (Table 1). The finding of KR1 was in close agreement with Mecheri sheep (Jeichitra and Rajendran, 2014) as  $13.07 \pm 0.06$  whereas higher estimate was reported in Nilagiri sheep as  $14.37 \pm 0.16$  and  $14.52 \pm 0.16$  in Sandyno sheep (Venkataramanan *et al.* 2016). The finding of KR2 was in close agreement with  $6.71 \pm 0.20$  in Harnali sheep (Bangar *et al.* 2020),  $6.19 \pm 0.04$  in Bharat Merino sheep (Mallick *et al.* 2019),  $6.83 \pm 0.11$  in Nellore sheep (Kumar *et al.* 2018) and  $6.10 \pm 0.07$  in Mecheri sheep (Jeichitra and Rajendran, 2014). The finding of KR3 was in close agreement with reports as  $3.82 \pm 0.22$  in Sandyno sheep and  $3.72 \pm 0.22$  in Nilagiri sheep (Venkataramanan *et al.* 2016). The finding of KR4 was higher than the value reported as  $3.72 \pm 0.21$  in Nilagiri sheep,  $3.80 \pm 0.22$  in Sandyno sheep (Venkataramanan *et al.* 2016) and  $3.34$  in Baluchi sheep (Pasandideh *et al.* 2018).

### Effect of sire

Sire had highly significant ( $P \leq 0.01$ ) effect on Kleiber ratio KR1, KR2 and KR3 whereas significant ( $P \leq 0.05$ ) on KR4. Similar results were reported in Pantja goat (Khadda *et al.* 2018) for KR1, KR2 and KR4.

### Effect of period of birth

Period of birth had highly significant ( $P \leq 0.01$ ) effect on KR1, KR2, KR3 and KR4. Similar results

**Table 1:** Least-squares means ( $\pm$ SE) of Kleiber ratio in Sonadi sheep

Effect	KR1	KR2	KR3	KR4
<b>Overall</b>	13.16 $\pm$ 0.30 (1060)	6.59 $\pm$ 0.33 (831)	4.04 $\pm$ 0.31 (630)	5.24 $\pm$ 0.35 (506)
<b>Sire</b>	**	**	**	*
<b>Period of birth</b>	**	**	**	**
P1 (2012-2013)	11.68 <sup>a</sup> $\pm$ 0.34 (318)	7.41 <sup>b</sup> $\pm$ 0.37 (257)	4.99 <sup>b</sup> $\pm$ 0.35 (210)	6.83 <sup>a</sup> $\pm$ 0.39 (184)
P2 (2014-2015)	11.65 <sup>a</sup> $\pm$ 0.34 (226)	7.83 <sup>b</sup> $\pm$ 0.37 (191)	4.82 <sup>b</sup> $\pm$ 0.35 (145)	5.50 <sup>a</sup> $\pm$ 0.39 (133)
P3 (2016-2017)	14.22 <sup>b</sup> $\pm$ 0.33 (292)	5.79 <sup>a</sup> $\pm$ 0.38 (196)	3.26 <sup>a</sup> $\pm$ 0.35 (143)	4.67 <sup>b</sup> $\pm$ 0.39 (121)
P4 (2018-2019)	15.10 <sup>c</sup> $\pm$ 0.34 (224)	5.32 <sup>a</sup> $\pm$ 0.37 (187)	3.11 <sup>a</sup> $\pm$ 0.35 (132)	3.95 <sup>a</sup> $\pm$ 0.42 (68)
<b>Season of birth</b>	NS	NS	NS	**
S1 (July-Oct.)	12.89 $\pm$ 0.31 (413)	6.53 $\pm$ 0.34 (354)	4.14 $\pm$ 0.31 (272)	5.03 <sup>a</sup> $\pm$ 0.36 (216)
S2 (Nov.-Feb.)	13.15 $\pm$ 0.30 (517)	6.90 $\pm$ 0.33 (384)	3.74 $\pm$ 0.31 (283)	4.82 <sup>a</sup> $\pm$ 0.36 (220)
S3 (March-June)	13.45 $\pm$ 0.39 (130)	6.33 $\pm$ 0.44 (93)	4.25 $\pm$ 0.40 (75)	5.87 <sup>b</sup> $\pm$ 0.44 (70)
<b>Sex of lamb</b>	NS	**	NS	NS
Male	13.31 $\pm$ 0.31 (509)	6.94 <sup>b</sup> $\pm$ 0.34 (400)	4.22 $\pm$ 0.32 (295)	5.37 $\pm$ 0.37 (236)
Female	13.02 $\pm$ 0.31 (551)	6.24 <sup>a</sup> $\pm$ 0.34 (431)	3.87 $\pm$ 0.32 (335)	5.11 $\pm$ 0.36 (270)
<b>Type of birth</b>	NS	NS	NS	*
Single	13.70 $\pm$ 0.11 (1031)	6.45 $\pm$ 0.12 (810)	4.31 $\pm$ 0.11 (614)	4.53 <sup>a</sup> $\pm$ 0.11 (495)
Twin	12.62 $\pm$ 0.58 (29)	6.72 $\pm$ 0.64 (21)	3.77 $\pm$ 0.60 (16)	5.95 <sup>b</sup> $\pm$ 0.69 (11)

\*\*=Highly significant ( $P \leq 0.01$ ), \*=Significant ( $P \leq 0.05$ ), NS= Non-significant ( $P > 0.05$ ) Figures in parentheses are the number of observations.

were reported as highly significant in Harnali sheep (Bangar *et al.* 2020), Bharat Merino sheep (Mallick *et al.* 2019) and Nellore sheep (Kumar *et al.* 2018) for KR1 and KR2. On the contrary, Nilagiri and Sandyno sheep (Venkataramanan *et al.* 2016) observed non-significant effect of period of birth on KR3 and KR4. Over the periods, there was increase the Kleiber ratio of Sonadi lambs, indicating production of more efficient animals for feed conversion.

#### Effect of season of birth

Season of birth had highly significant ( $P \leq 0.01$ ) effect on KR4 whereas non-significant effect on KR2, KR3 and KR4. Similar results were reported as non-significant effect in Mehsana goat (Gupta *et al.* 2016) and Pantja goat (Khadda *et al.* 2018) for KR1. Similar results were reported as non-significant effect in Harnali sheep (Bangar *et al.* 2020) and Pantja goat (Khadda *et al.* 2018) for KR2. On the contrary, results were reported in Pantja goat (Khadda *et al.* 2018) on KR3 and KR4. There was increase the Kleiber ratio of Sonadi lambs during summer season, indicating production of more efficient animals for feed conversion.

#### Effect of sex of lamb

Sex of lamb had highly significant ( $P \leq 0.01$ ) effect on KR2 whereas non-significant effect on KR1, KR3 and KR4. Similar results were reported as non-significant effect in Deccani sheep (Bangar *et al.* 2018) and Nellore sheep (Kumar *et al.* 2016) for KR1. Similar results were reported as highly significant in Harnali sheep (Bangar *et al.* 2020), Bharat Merino sheep (Mallick *et al.* 2019) and Nilagiri and Sandyno sheep (Venkataramanan *et al.* 2016) for KR2. Similar results were reported as non-significant in Pantja goat (Khadda *et al.* 2018) for KR3 and KR4. Males usually have more feed conversion than females at all stages of growth.

#### Effect of type of birth

Type of birth had significant ( $P \leq 0.05$ ) effect on KR4 whereas non-significant effect on KR1, KR2 and KR3. Similar result was reported as non-significant effect in Raeini Cashmere goat (Mokhtari *et al.* 2019) for KR1 and KR2. Similar results were reported as non-significant in Nilagiri and Sandyno sheep (Venkataramanan *et al.* 2016) and Pantja goat (Khadda *et al.* 2018) for KR3. On the contrary, reported non-significant effect of type of birth in

**Table 2:** Estimates of heritability (diagonal), genetic (above diagonal) and phenotypic (below diagonal) correlations along with standard error among Kleiber ratio traits in Sonadi sheep

Traits	KR1	KR2	KR3	KR4
KR1	0.82±0.18	-0.76±0.28	-0.79±0.19	-1.00±0.31
KR2	-0.52	0.38±0.11	0.78±0.28	0.86±0.39
KR3	-0.38	-0.05	0.51±0.15	0.69±0.32
KR4	-0.33	-0.03	-0.04	0.23±0.13

Pantja goat (Khadda *et al.* 2018) and Nilagiri and Sandyno sheep (Venkataramanan *et al.* 2016) for KR4. Single lamb had higher Kleiber ratio than twin lambs because of compete feeding of mother milk in twin lambs.

### Genetic parameters

Estimates of heritability for Kleiber ratio KR1, KR2, KR3 and KR4 were observed as 0.82±0.18, 0.38±0.11, 0.51±0.15 and 0.23±0.13, respectively (Table 2). The estimates of heritability for KR1 and KR2 were higher than reports as 0.38±0.11 and 0.02±0.06 in Harnali sheep (Bangar *et al.* 2020), 0.17±0.02 and 0.17±0.01 in Bharat Merino sheep (Mallick *et al.* 2019) and 0.106±0.03 and 0.106±0.03 in Nilagiri and Sandyno sheep (Venkataramanan *et al.* 2016), respectively. The present finding of heritability for KR3 and KR4 were higher than reports as 0.22±0.08 and 0.04±0.03 in Pantja goat (Khadda *et al.* 2018) and 0.035±0.03 and 0.048±0.03 in Nilagiri and Sandyno sheep (Venkataramanan *et al.* 2016), respectively. The heritability estimates provides a scope that individual selection may be used for selection in case of non-availability of other information.

The estimates of genetic correlation among Kleiber ratio traits were high in magnitude and ranged from -0.69 (KR3-KR4) to -1.00 (KR1-KR4). The present finding of genetic correlation among Kleiber ratio for KR1-KR2, KR1-KR3 and KR1-KR4 were higher than reports as -0.285, -0.572 and -0.817 in Nilagiri sheep (Venkataramanan *et al.* 2016) and 0.41, -0.68 and -0.22 in Pantja goat (Khadda *et al.* 2018), respectively. Positive genetic correlations between Kleiber ratio traits indicated that the selection of one trait can improve other traits simultaneously. Negative genetic correlation between Kleiber ratios were estimated due to maternal compensatory effect.

The estimates of phenotypic correlation among Kleiber ratio traits were negative and magnitude is

very low to high and ranged from -0.03 (KR2-KR4) to -0.52 (KR1-KR2). The present finding of phenotypic correlation for KR1-KR2, KR1-KR3 and KR1-KR4 were higher than reports as -0.358, -0.077 and -0.167 in Nilagiri sheep (Venkataramanan *et al.* 2016) and -0.43, -0.30 and -0.01 in Pantja goat (Khadda *et al.* 2018), respectively. Negative phenotypic correlation between Kleiber ratio traits were estimated due to maternal compensatory effect.

### CONCLUSION

Assessment of non-genetic factors plays an important role to formulate effective breeding programme for improvement of growth traits of Sonadi sheep and measure to be taken for standardizing the management of flock for suitable production. More lambing should be planned in conducive season such as summer season for maximizing the growth rate and survivability of lambs. Lambs showing multiple birth must be given preference in breeding programme to improve prolificacy of the flock. Kleiber ratio is more important trait for efficiency of feed utilization. Medium to high heritability of Kleiber ratio offer a scope for individual selection for further genetic improvement.

### ACKNOWLEDGEMENTS

The author is thankful to Mega Sheep Seed Project (MSSP) on Sonadi Sheep, Vallabh Nagar, Udaipur, for providing data and all the necessary facilities to conduct the research.

### REFERENCES

- Arthur, P.F., Renand, G. and Krauss, D. 2001. Genetic and phenotypic relationships among different measures of growth and feed efficiency in young Charolais bulls. *Livest. Prod. Sci.*, **68**(2-3): 131-139.
- Bangar, Y.C., Lawar, V.S., Nimbalkar, C.A., Shinde, O.V. and Nimase, R.G. 2018. Heritability estimates for average daily



- gain and Kleiber ratio in Deccani sheep. *Indian J. Small Rumin.*, **24**(1): 18-21.
- Bangar, Y.C., Magotra, A. and Yadav, A.S. 2020. Estimates of covariance components and genetic parameters for growth, average daily gain and Kleiber ratio in Harnali sheep. *Trop. Anim. Health Prod.*, **52**: 1-6.
- Eskandarinasab, M., Ghafouri-Kesbi, F. and Abbasi, M.A. 2010. Different models for evaluation of growth traits and Kleiber ratio in an experimental flock of Iranian fat-tailed Afshari sheep. *J. Anim. Breed. Genet.*, **127**(1): 26-33.
- Gupta, J.P., Pandey, D.P., Panchasara, H.H. and Shah, R.R. 2016. Factors affecting pre- and post- weaning Kleiber ratios and genetic parameters in Mehsana goats. *Indian J. Small Rumin.*, **22**(1): 100-102.
- Harvey, W.R. 1990. User's guide for LSMLMW and MIXMDL PC-2 version. Mixed model least squares and maximum likelihood computer program, Ohio State University, Columbus, Ohio, U.S.A.
- Jeichitra, V. and Rajendran, R. 2014. Kleiber Ratio in Mecheri Sheep. *Indian Vet. J.*, **91**(07): 17-19.
- Khadra, B.S., Singh, B., Singh, D.V., Singh, S.K., Singh, C.B. and Singh, J.L. 2018. Genetic and non-genetic parameters of Kleiber ratio in Pantja goat under field conditions of Tarai region of Uttarakhand. *Indian J. Anim. Sci.*, **88**(12): 1370-1373.
- Kleiber, M. 1947. Body size and metabolic rate. *Physiol Rev.*, **27**(4): 511-541.
- Kramer, C.Y. 1956. Extension of multiple range tests to group means with unequal numbers of replications. *Biometrics*, **12**: 307-310.
- Kumar, D., Prakash, M.G., Gupta, B.R., Raghunandan, T.R. and Chandra, A.S. 2017. Average daily gain and Kleiber Ratio in Deccani Sheep. *J. Pharma Innov.*, **6**: 194-197.
- Kumar, I.S., Gangaraju, G., Kumar, C.V. and Nath, S. 2018. Genetic parameters for growth rate and Kleiber ratios of Nellore sheep. *Indian J. Anim. Res.*, **52**(10): 1405-1408.
- Livestock Census. 2019. Basic animal husbandary statistics. Department of Animal Husbandry, Dairying and Fisheries, Government of India. (<http://www.dahd.nic.in>).
- Mallick, P.K., Chauhan, I.S., Gowane, G.R., Thirumurgan, P., Murali, G. and Kumar, A. 2019. Average daily gain and Kleiber ratio in Bharat merino sheep. *Indian J. Small Rumin.*, **25**(2): 151-155.
- Mokhtari, M.S., Razmkabir, M., Ghiasi, H. and Mohammadi, Y. 2019. Genetic evaluation of growth rate and efficiency related traits in Raeini Cashmere goat. *Iran. J. Appl. Anim Sci.*, **9**(2): 275-282.
- Pasandideh, M., Rahimi-Mianji, G. and Gholizadeh, M. 2018. A genome scan for quantitative trait loci affecting average daily gain and Kleiber ratio in Baluchi Sheep. *J. Genet.*, **97**(2): 493-503.
- Venkataramanan, R., Subramanian, A., Sivaselvam, S.N., Sivakumar, T. and Sreekumar, C. 2016. Genetic parameters for Kleiber ratio and its relation to other body weight traits in Nilagiri and Sandyno sheep. *Indian J. Anim. Sci.*, **86**(5): 559-563.

